

9-2015

Challenging the efficacy of different dental bleaching protocols

A. Koniaris

Vasileios Margaritis
Walden University

Follow this and additional works at: http://scholarworks.waldenu.edu/shs_pubs



Part of the [Dentistry Commons](#)

This Conference Proceeding is brought to you for free and open access by the College of Health Sciences at ScholarWorks. It has been accepted for inclusion in School of Health Sciences Publications by an authorized administrator of ScholarWorks. For more information, please contact ScholarWorks@waldenu.edu.

A. Koniaris*, V. Margaritis, M. Antoniadou and A. Kakaboura

OPERATIVE DENTISTRY DEPARTMENT, DENTAL SCHOOL,
UNIVERSITY OF ATHENS, GREECE

PURPOSE. Investigation of the *in vitro* efficiency of four dental bleaching protocols on the color change of teeth, stained by a black tea solution.

MATERIALS AND METHODS. One hundred intact, extracted, human incisors were randomly divided (n=4x25/group) and underwent black tea staining {2gr black tea (M&S Fairtrade Earl Grey) at 100ml boiled distilled water}, for seven days in room temperature (Fig.1).

The bleaching protocols applied per group were¹⁻⁴:

- A. 14 days x 120 min/day 10% CP (Opalescence PF 10%, Ultradent) with trays (bleaching at home-BH),
- B. 2 times x (3 appl/time x 15min) 40% HP, (Opalescence Boost 40%, Ultradent) (bleaching in office-BO1),
- C. 3 times x (3 appl/time x 15min) 40% HP, (Opalescence Boost 40%, Ultradent) (bleaching in office-BO2) and,
- D. 14 days x 120 min/day 10% CP with trays and 2 times x (3appl/time x 15 min) 40% HP (combined BH-BO1).




Fig. 1. Materials and methods of the study: a. Tooth numbering, b. Stained natural teeth fixed on frasaco and impression taking, c. Dental cast for splint preparation d. In-office whitening for groups B and C, e. Colorimeter.

The color measurements were conducted at baseline (t_0) and after staining (t_1), bleaching application (t_2), 90 (t_3) and 180 (t_4) days, with a colorimeter (Dr Lance Micro Colour, Braive Instruments) in CIEL*a*b* system. ΔE values were calculated relative to t_0 and analyzed by three-way ANOVA, Mann-Whitney U and Wilcoxon tests ($\alpha=0.01$).

RESULTS. There were no group-dependenced differences for ΔE values and L^* was significantly different at all times for all groups.

All groups presented augmenting L^* values from t_1 to t_2 and diminishing from t_2 to t_4 (Fig.2). BH had significant differences in L^* values at all-times. At t_3 and t_4 the lowest L^* and the highest ΔE were reported for BH while the requested combination of a high L^* and a low ΔE was reported to be best for BH-BO1 (Table 1).




Fig. 2. Graphs presenting L^* , a^* , b^* and ΔE variation over time

Whitening protocols	ΔE			
	t_1 (baseline)	t_2 (after staining)	t_3 (90 days)	t_4 (180 days)
Group 1 (BH)	9,53 ($\pm 5,61$) ^{1,2,3}	5,20 ($\pm 4,05$) ^{1,4,5}	12,59 ($\pm 6,43$) ^{2,4,6}	15,22 ($\pm 6,47$) ^{3,5,6}
Group 2 (BO1)	10,36 ($\pm 4,27$) ¹	5,87 ($\pm 5,04$) ^{1,2,3}	9,48 ($\pm 3,17$) ^{2,4}	10,77 ($\pm 3,37$) ^{3,4}
Group 3 (BO2)	11,67 ($\pm 5,93$) ¹	7,31 ($\pm 4,66$) ^{1,2,3}	11,19 ($\pm 6,63$) ²	12,43 ($\pm 5,94$) ³
Group 4 (BH-BO1)	12,41 ($\pm 5,31$) ^{1,2}	4,88 ($\pm 2,89$) ^{1,3,4}	9,84 ($\pm 4,55$) ^{2,3,5}	11,71 ($\pm 4,61$) ^{4,5}

Table 1. Mean ΔE values for all groups at all times. Same numbers within the rows for each parameter (L^* , a^* , b^*) indicate statistically significant differences between time points for each group/row, Friedman test and post-hoc Wilcoxon Signed Ranks test, $p < 0.01$.

CONCLUSIONS. In-office, at-home and combined bleaching protocols proved to be equally efficient although several differences were detected concerning color coordinates. All protocols tested provided significant color change in stained teeth just after bleaching within the same group. BH had the highest relapse, while BH-BO1 reported better bleaching efficiency at 6 months.

Literature

1. Dietschi D, Benbachir N, Krejci I. In vitro colorimetric evaluation of the efficacy of home bleaching and over-the-counter bleaching products. *Quintessence Int* 2010; 41(6): 505-516.
2. Scaminaci Russo D, et al. "Color stability of bleached teeth over time: an in vitro study." *Eur J Esthet Dent* 2010; 5(3): 300-310.
3. Giachetti L, et al. "A randomized clinical trial comparing at-home and in-office tooth whitening techniques: A nine-month follow-up." *J Am Dent Assoc* 2010;141(11): 1357-1364.
4. Dawson PF, et al. "A clinical study comparing the efficacy and sensitivity of home vs combined whitening." *Oper Dent* 2011; 36(5): 460-466.