Walden University
ScholarWorks

Walden Dissertations and Doctoral Studies

Walden Dissertations and Doctoral Studies Collection

8-29-2025

Staff Education to Reduce Alarm Fatigue and Enhance Patient Safety Through Nurse Education on Telemetry Optimization

Casey Alan Metcalf
Walden University

Follow this and additional works at: https://scholarworks.waldenu.edu/dissertations

 Part of the Nursing Commons

This Dissertation is brought to you for free and open access by the Walden Dissertations and Doctoral Studies Collection at ScholarWorks. It has been accepted for inclusion in Walden Dissertations and Doctoral Studies by an authorized administrator of ScholarWorks. For more information, please contact ScholarWorks@waldenu.edu.

Walden University

College of Nursing

This is to certify that the doctoral study by

Casey Metcalf

has been found to be complete and satisfactory in all respects,
and that any and all revisions required by
the review committee have been made.

Review Committee
Dr. Maria Revell, Committee Chairperson, Nursing Faculty
Dr. Diane Whitehead, Committee Member, Nursing Faculty

Chief Academic Officer and Provost
Sue Subocz, Ph.D.

Walden University
2025

Staff Education to Reduce Alarm Fatigue and Enhance Patient Safety Through Nurse Education on Telemetry Optimization

by

Casey Metcalf

MS, Walden University, 2022

BS, Jacksonville State University, 2010

Executive Summary Submitted in Partial Fulfillment

of the Requirements for the Degree of

Doctor of Nursing Practice

Walden University

November 2025

Summary

This project was an evidence-based staff education initiative to reduce alarm fatigue and enhance patient safety through telemetry optimization in an acute care setting. The core issue involved the overuse of continuous cardiac monitoring, which contributed to nonactionable alarms, alarm fatigue, workflow inefficiencies, delayed throughput, and increased costs. The project was a structured intervention delivered to 171 telemetry nurses across three inpatient units. Pretests and posttests measured changes in knowledge, intent, and practice. Results showed a 22.9% increase in knowledge, a 92% agreement regarding intention to follow discontinuation protocols, and a 10% overall reduction in practice of telemetry use on two of the three units. Likert scores for implementation intention demonstrated improvement in scores and overall percent for all units (1W, 3.41 to 4.26, 25%; 2W, 4.26 to 4.48, 5.2%; 3W, 4.14 to 4.25, 2.6%). The mean post-30-day Likert score was 3/5, indicating overall neutrality in sustaining project practices. However, participants reported higher confidence in discontinuing monitoring (3.4/5), perceived effectiveness in reducing alarm fatigue (3.2/5), and improved patient safety through reduced false alarms and better transitions (3.5/5). Implementing equitable, evidence-based monitoring practices can standardize care delivery, foster psychological safety in the workplace, and improve nurse well-being, retention, and health equity. This project included an educational design that has the potential to enhance patient telemetry management, reduce care gaps, and promote positive social change by mitigating alarm fatigue and improving patient safety in clinical settings.

Background

The background of this project included a persistent practice gap in the overutilization of telemetry monitoring, which led to alarm fatigue in acute care settings. Although telemetry is a critical tool for monitoring cardiac function, inappropriate or prolonged use has been associated with increased nonactionable alarms, nurse desensitization, delayed response times, and higher operational costs (Sandau et al., 2020). The lack of standardized protocols for initiating and discontinuing telemetry, limited nurse education on alarm management, and underutilization of clinical decision-support tools contribute to compromised patient safety and workflow inefficiencies (Lehet et al., 2023). These challenges provided the foundation for the identified practice gap and the need for targeted intervention.

The practice-focused question guiding the project addressed whether structured nursing education in acute care inpatient units reduces alarm fatigue, improves telemetry discontinuation rates, and enhances telemetry transitions over a 30-day period. The purpose of the project was to implement and evaluate an educational intervention that would optimize telemetry use, reduce alarm fatigue, and align nursing practices with evidence-based guidelines. This focus addressed persistent barriers in acute care where poor alarm management, including frequent nonactionable alerts, undermines clinical vigilance, prolongs response times, and elevates patient safety risks (Chen et al., 2023). Despite the availability of national practice standards, many telemetry units operate without standardized discontinuation protocols or consistent staff training. Structured education and targeted process improvements are crucial for shifting practice toward evidence-based monitoring, reducing unnecessary utilization, and enhancing care

coordination (Pendharkar et al., 2020). Educational interventions that emphasize clinical judgment and adherence to protocols have demonstrated success in reducing inappropriate use of telemetry and hospital costs (Rodriguez et al., 2024). This project served as the foundation for reducing telemetry overuse, enhancing workflow efficiency, and supporting safer, more cost-effective care.

A synthesis of the evidence confirmed that implementing structured nursing education on telemetry optimization is feasible and supported by the literature. This intervention addresses critical concerns of alarm fatigue, inefficient utilization of telemetry, and delays in patient throughput. Evidence consistently showed that structured education on integrating standardized telemetry protocols enhances clinical decision making, reduces nonactionable alarms, and improves patient safety and workflow efficiency (Rodriguez et al., 2024; Sandau et al., 2020).

The literature supporting this project encompassed a diverse and rigorous body of research. Bi et al. (2020) conducted a randomized controlled trial demonstrating that alarm management training significantly decreased alarm fatigue and improved nurses' responses to clinically relevant alerts. Chen et al. (2023), through a systematic review, identified key modifiable contributors to alarm fatigue, including lack of standardized protocols and insufficient staff training, further underscoring the need for targeted educational interventions. Pendharkar et al. (2020) employed a quasi-experimental design and demonstrated that implementing American Heart Association (AHA) telemetry guidelines resulted in substantial reductions in inappropriate monitoring and improved resource utilization efficiency.

Additionally, Rodriguez et al. (2024) conducted a quality improvement project in which structured education on discontinuing telemetry for providers led to greater protocol adherence and meaningful cost savings. The significance of education in promoting safe telemetry practices was underscored by Sandau et al. (2020), who emphasized the importance of updated electrocardiographic monitoring standards and called for routine reassessment and staff education to ensure appropriate use. Bergstedt et al. (2024) evaluated a clinical decision support tool aligned with AHA standards, highlighting its positive influence on promoting evidence-based monitoring and reducing unnecessary telemetry. Moreover, Duffy et al. (2020) demonstrated that empowering nurses and residents through telemetry stewardship initiatives can lead to sustainable reductions in overmonitoring while maintaining clinical safety. Collectively, these findings affirmed that a structured, evidence-based educational intervention focused on reducing unnecessary telemetry use is not only justified but also essential for addressing alarm fatigue in acute care environments. Educating nurses on appropriate telemetry indications and discontinuation protocols enhances clinical judgment, promotes adherence to evidence-based guidelines, and contributes to reducing nonactionable alarms, thereby improving patient safety, workflow efficiency, and nurse well-being.

Staff Education Project Development

The project was implemented in an acute care hospital setting and involved 171 frontline registered nurses working across three inpatient telemetry units. These participants were responsible for cardiac monitoring and represented the primary target audience for the structured educational intervention. The organizational readiness assessment revealed that the hospital was moderately prepared for change, with

leadership demonstrating a supportive culture and a strong commitment to improving patient care. Although some resistance was noted among staff primarily due to uncertainty about protocol changes and concerns about workload, leadership's investment in education, resource allocation, and ongoing support created a foundation for successful implementation. A strategic stakeholder communication plan was developed to raise awareness, promote adherence to telemetry guidelines, and engage frontline nurses, interdisciplinary partners, and leadership through meetings, briefings, policy updates, and training sessions. These efforts ensured alignment across roles and sustained engagement throughout the intervention.

A strengths, weaknesses, opportunities, and threats analysis identified key institutional assets, including adherence to evidence-based guidelines, capacity for resource optimization, and a culture that values clinical education. Challenges such as change fatigue, initial implementation costs, and limited awareness of telemetry protocols were acknowledged and proactively addressed. Growth opportunities included leveraging advanced monitoring technologies, strengthening telemetry-related training, and reducing unnecessary utilization. By capitalizing on internal strengths and mitigating known risks, the organization was well positioned to sustain this educational initiative.

The intervention consisted of targeted educational sessions with a PowerPoint that addressed telemetry guidelines, alarm fatigue, and protocols for discontinuation (see Appendix A). Twelve sessions over 2 weeks provided flexible opportunities for all telemetry staff. There were six sessions each for the day and night shift. Makeup sessions were provided for staff who were off duty or unavailable. Each session was 45 to 60 minutes in length and included a structured presentation, case-based discussions, and

distribution of a telemetry optimization toolkit. Content focused on five key educational objectives: (a) reducing false alarms, (b) mitigating alarm fatigue, (c) improving telemetry transitions and discharge efficiency, (d) increasing intent to follow discontinuation protocols, and (e) decreasing overall telemetry use.

The evaluation process included quantitative methods to assess the effectiveness of educational intervention. Nurse knowledge was evaluated using a 20-question multiple choice pretest (see Appendix B) and posttest (see Appendix C), while self-reported intent to follow telemetry discontinuation protocols was measured using an eight-item Likert scale. Participant satisfaction was assessed through an end-of-session survey, which included items evaluating content clarity, relevance to clinical practice, and achievement of learning objectives (see Appendix D). The project aimed for a minimum 20% increase in test scores and 90% positive response rate regarding nurses' intent to follow telemetry protocols.

To determine the sustained impact of the intervention, I conducted a 30-day follow-up survey to evaluate perceptions of alarm fatigue, confidence in applying telemetry guidelines, and perceived improvements in patient safety and workflow efficiency (see Appendix E). Organizational telemetry data were collected to assess baseline and postintervention telemetry order counts and average monitoring duration to achieve at least a 10% reduction in use. Outcome metrics included telemetry order counts and average monitoring duration, comparing 30 days preceding and following the intervention. This multimethod approach provided a comprehensive understanding of the intervention's influence on nursing practice, staff engagement, and system-level outcomes.

Results

A total of 171 RNs participated in the educational intervention. Of these, 61 were from Unit 1W (35.7%), 54 from Unit 2W (31.7%), and 56 from Unit 3W (32.6%). Pretest scores were 72% in Unit 1W, 67% in Unit 2W, and 70% in Unit 3W. Posttest scores increased to 86% in Unit 1W, 84% in Unit 2W, and 86% in Unit 3W. Pretest scores ranged from 35% to 100% across all units, while posttest scores ranged from 60% to 100%, indicating consistent improvement across the entire performance range. The mean score improved from 70% preintervention to 86.1% postintervention, representing a 22.9% percentage change. All units met the objective of increasing the 20% score.

Baseline telemetry utilization was assessed using organizational data, which captured the average duration of telemetry monitoring across three inpatient units. Unit 1W, a surgical step-down unit, achieved the most substantial reduction, decreasing average telemetry use by 18% from 90 hours to 73.8 hours. Similarly, Unit 2W, which serves a medical oncology population, demonstrated a 17% reduction, lowering average telemetry duration from 67 hours to 55.6 hours. Units 1W and 2W met the 10% reduction in telemetry usage objective. In contrast, Unit 3W, which primarily admits patients with neurological conditions and stroke, showed a 3% reduction from 72 hours to 70 hours, which did not meet the 10% objective (see Figure 1).

The educational intervention met its objective of increasing nurses' intent to follow telemetry discontinuation protocols when clinically appropriate. Based on postsession Likert scale data, 92% of participants reported a score of 4 or 5 (*agree* or *strongly agree*), exceeding the target threshold of 90%.

Figure 1

Telemetry Utilization



The mean intention to implement pretest Likert score for Unit 1W was 3.41, increasing to 4.26 postintervention, a 25.0% improvement. Unit 2W maintained a high baseline with a pretest mean of 4.26, which rose to 4.48, a 5.2% improvement, while Unit 3W improved from 4.14 to 4.25, a 2.6% improvement. These gains reflect improved awareness and reinforced confidence in protocol adherence. The immediate posttest responses indicated high levels of understanding and engagement with the intervention content across all telemetry units.

Based on the 30-day postproject survey, the follow-up Likert scores had a mean of 3.0, with scores ranging from 1 to 5. The mean identified neutrality among participants regarding sustained confidence in telemetry protocol adherence and the perceived impact of the intervention on clinical practice. Participants expressed confidence in applying AHA guidelines (mean score 3.3) and determining when telemetry was no longer

necessary (mean score 3.4), aligning with the measurable and achievable components of the goal. Responses confirmed that the project improved patient safety by reducing false alarms and enhancing transitions (mean score 3.5).

These outcomes reflect meaningful system-level improvements, including enhanced nursing knowledge, increased adherence to telemetry discontinuation protocols, and improved efficiency in monitoring practices. By reducing unnecessary telemetry and aligning monitoring with evidence-based guidelines, the intervention supported patient safety and optimized resource use. The intervention contributed to reduced alarm fatigue, which is a factor that influences nurse workload, throughput, and clinical decision making. The balanced unit-level engagement fostered a shared sense of accountability.

Despite the overall success, several limitations affected the results. Unit 3W demonstrated only a 3% reduction in telemetry duration due to regulatory protocols requiring extended cardiac monitoring in stroke patients, which limited opportunities for reduction and skewed unit-level impact comparisons. Additionally, reliance on self-reported intent and satisfaction may have introduced bias, and the project's short-term evaluation period limited assessment of sustainability and patient-level outcomes.

This project addressed a nationally recognized issue in acute care: alarm fatigue and inappropriate telemetry use. The project demonstrated that targeted nursing education can lead to measurable improvements in knowledge, practice, and resource efficiency. By aligning telemetry practices with AHA guidelines, the intervention model can be transferred to similar inpatient settings that face high volumes of nonactionable alarms and prolonged monitoring periods. Broader adoption of this approach may promote

equitable, evidence-informed care and foster safer clinical environments across diverse institutions.

Conclusions

The project demonstrated measurable improvements in organizational performance, clinical decision making, and protocol adherence. Key outcomes included a 22.9% increase in knowledge, 92% agreement to follow telemetry discontinuation protocols postintervention, and a 13% reduction in monitoring duration. These results reflect enhanced nursing competency in managing alarm fatigue and applying evidence-based monitoring, contributing to improved patient safety, workflow efficiency, and resource stewardship. The intervention's alignment with AHA guidelines, institutional quality goals, and national standards supports its scalability and sustainability. The intervention is low-risk, cost-effective, and practical for long-term adoption.

To sustain progress, recommendations include ongoing audit and feedback, telemetry alerts, and annual refresher training. Future evaluation should assess long-term effects on patient outcomes, staff satisfaction, and transfer efficiency. Given that Unit 3W's telemetry use was impacted by regulatory requirements, future efforts should involve collaboration with case management and compliance teams.

Broader implications include strengthening nursing practice and promoting social change. By empowering nurses to make informed telemetry decisions, the project supports clinical autonomy, consistency, and equitable care. This fosters a psychologically safe environment and advances nurse well-being, retention, and health equity.

References

Bergstedt, A., Hilliard, B., Alabsi, S., Usher, M. G., Peters, M., Grace, J., Melton, G. B., Beebe, T. J., & Pestka, D. L. (2024). Evaluation of a clinical decision support tool to guide adoption of the American Heart Association telemetry monitoring practice standards. *Journal of the American Heart Association*, *13*(9), Article e031523. https://doi.org/10.1161/JAHA.123.031523

Bi, J., Yin, X., Li, H., Gao, R., Zhang, Q., Zhong, T., Zen, T., Guan, B., & Li, Z. (2020). Effects of monitor alarm management training on nurses' alarm fatigue: A randomized controlled trial. *Journal of Clinical Nursing*, *29*(21–22), 4203–4216. https://doi.org/10.1111/jocn.15452

Chen, S., Hui, Z., Su, W., Ruxin, J., Huiyi, Z., Hongmei, Z., & Hongyan, Z. (2023). Determinants of medical equipment alarm fatigue in practicing nurses: A systematic review. *SAGE Open Nursing*, *9*, 1–10. https://doi.org/10.1177/23779608231207227

Duffy, E., Niessen, T., Perrin, K., Apfel, A., Bertram, A., Keller, S. C., Feldman, L. S., & Pahwa, A. K. (2020). Empowering nurses and residents to improve telemetry stewardship in the academic care setting. *Journal of Evaluation in Clinical Practice*, *27*(5), 1154–1158. https://doi.org/10.1111/jep.13470

Lehet, C. R., Lopez, J. A., Frank, R. J., & Cvach, M. (2023). Technological intervention to improve alarm management in acute care telemetry units. *Biomedical Instrumentation & Technology*, *57*(2), 67–75. https://doi.org/10.2345/0899-8205-57.2.67

Pendharkar, S. S., Barry, I. B., Patibandla, S., Leung, T. I., Gupta, A., Lin, A. N., & Gasperino, J. (2020). AHA telemetry guidelines improve telemetry utilization in the inpatient setting. *The American Journal of Managed Care*, *26*(11), 476–481. https://doi.org/10.37765/ajmc.2020.88525

Rodriguez, C., Bianco, N., Bucco, T., Collum, K., O'Neill, S. P., & David, D. (2024). Telemetry discontinuation education for nurse practitioners decreases hospital costs: A quality-improvement project. *Journal of the American Association of Nurse Practitioners*, *36*(10), 576–585. https://doi.org/10.1097/JXX.0000000000001062

Sandau, K. E., Funk, M., Auerbach, A., Barsness, G. W., Blum, K., Cvach, M., Lampert, R., May, J. L., McDaniel, G. M., Perez, M. V., Sendelbach, S., Sommargren, C. E., & Wang, P. J. (2020). Update to practice standards for electrocardiographic monitoring in hospital settings: A scientific statement from the American Heart Association. *Circulation*, *141*(3), Article e55e70. https://doi.org/10.1161/CIR.0000000000000754

Appendix A: Education PowerPoint

Objectives

Improve	Improve nurse knowledge of false alarms and telemetry transitions.
Reduce	Reduce alarm fatigue and enhance patient safety.
Increase	Increase adherence to telemetry discontinuation protocols.



Introduction to Telemetry Optimization

- Telemetry: Continuous monitoring of a patient's heart rhythm.
- Essential for early detection of cardiac events.
- Overuse or mismanagement can lead to alarm fatigue.

Alarm Fatigue and Its Consequence

Definition: Desensitization to alarms due to high frequency.

- **Causes:**
  - **High frequency of false alarms.**
  - **Lack of standardized protocols.**
  - **Environmental noise.**
- **Impact:**
  - **Delayed response to critical events.**
  - **Increased stress and burnout.**



Types of Alarms

☑ Actionable Alarms

- Require immediate clinical intervention.
- Examples:
  - Cardiac arrest (asystole).
  - Respiratory failure.
  - Life-threatening arrhythmias (e.g., ventricular tachycardia).

✕ Non-Actionable Alarms

- Do not require clinical intervention.
- Examples:
  - Lead displacement.
  - Low battery warning.
  - Motion artifacts (e.g., patient adjusting position).

Impact of False Alarms on Patient Safety

- Excessive false alarms → alarm fatigue.
- Leads to desensitization → missed critical alarms.
- Consequences: Delayed responses and compromised patient safety.

Strategies to Reduce Alarm Fatigue

Customize	Customize alarm settings based on clinical needs.
Review and adjust	Regularly review and adjust telemetry orders.
Educate	Educate staff on alarm management.
Encourage	Encourage interdisciplinary collaboration.

AHA Guidelines

The American Heart Association (AHA) provides comprehensive guidelines for electrocardiographic (ECG) monitoring in hospital settings, aiming to optimize patient care and resource utilization. These guidelines address appropriate indications for telemetry use, duration of monitoring, and implementation strategies (Sandau et al., 2020).



AHA Guidelines

Clinical indications for telemetry:	Arrhythmias, acute coronary syndrome, heart failure. High-risk patients needing continuous monitoring.
Not recommended for:	Physician preference alone. Low-risk, stable patients.

AHA Guidelines

Indications for Telemetry Monitoring:

- Patients with acute coronary syndromes, significant arrhythmias, or conditions predisposing to arrhythmias are recommended for continuous ECG monitoring.
- Routine telemetry for low-risk patients without cardiac conditions is generally not recommended.

Duration of Monitoring:

- Continuous ECG monitoring should be maintained as long as clinically necessary, with regular assessments to determine the ongoing need.

AHA Guidelines

Alarm Management:	• Implement strategies to reduce non-actionable alarms, such as customizing alarm parameters and regularly maintaining monitoring equipment to mitigate alarm fatigue.
Duration of Monitoring:	• Provide ongoing education and training for healthcare professionals on appropriate telemetry use, alarm management, and response protocols to ensure adherence to guidelines.

Telemetry Transitions and Patient Discharge

- Prolonged telemetry use can:
  - Delay discharge.
  - Lead to unnecessary interventions.
- Best Practices:
  - Assess the need for telemetry daily.
  - Follow AHA guidelines for telemetry discontinuation.

Environmental Factors and Noise Pollution

- Noise pollution from constant alarms increases stress and cognitive overload.
- Strategies to reduce environmental noise:
  - Adjust alarm thresholds.
  - Use sound-absorbing materials.
  - Reduce background noise from equipment and staff communication.

Best Practices for Telemetry Use

- Guidelines for appropriate monitoring.
- Discontinuation protocols.
- Nurse-led decision-making.
- Data-driven adjustments.

Role of Clinical Decision Support Tools

- Tools assist in identifying appropriate telemetry use.
- Improve adherence to discontinuation protocols and standards.
- Reduce alarm frequency and increase nurse responsiveness.

Examples:

- A Nurse-Driven Telemetry Discontinuation Protocol
- EHR-integrated tool that required providers to justify telemetry orders by selecting from approved indications (e.g., arrhythmia, ischemia).

Nurse-Led Protocols and Training

Nurse-led telemetry discontinuation protocols:

Empower nurses to make decisions.
Improve staff confidence.
Increase adherence to guidelines.

Regular training enhances protocol compliance.

Conclusion

EFFECTIVE TELEMETRY USE IMPROVES PATIENT SAFETY.

REDUCING ALARM FATIGUE ENHANCES CLINICAL OUTCOMES.

NURSE EDUCATION AND ADHERENCE TO PROTOCOLS ARE KEY TO SUCCESS.

Questions and Discussion

Case Study: Reducing Alarm Fatigue and Enhancing Patient Safety

Background: A 58-year-old male patient is admitted to the telemetry unit following a myocardial infarction. He is placed on continuous telemetry monitoring to track cardiac activity. After stabilization, he begins experiencing frequent, non -critical alarm activations. The nursing staff reports an average of 40 alarms per hour, most of which are identified as false alarms due to patient movement and low battery warnings.

Situation: The telemetry unit has experienced increased false alarms, leading to alarm fatigue among the nursing staff. Multiple nurses have reported becoming desensitized to the sound of alarms, resulting in delayed responses to genuine alerts. Additionally, prolonged telemetry monitoring for stable patients has contributed to delayed discharge and increased workload for the staff

Case Study: Reducing Alarm Fatigue and Enhancing Patient Safety



Case Study: Reducing Alarm Fatigue and Enhancing Patient Safety

Intervention:

Education: Nurses participated in a telemetry optimization training program focused on:

- Identifying appropriate use of telemetry monitoring.
- Customizing alarm settings to minimize false alarms.
- Following the AHA guidelines for telemetry discontinuation.

Policy Update: Implemented a nurse-managed telemetry discontinuation protocol.

Monitoring: Collected baseline and post-intervention data on telemetry usage, alarm response times, and patient discharge rates.

Case Study: Reducing Alarm Fatigue and Enhancing Patient Safety

Discussion Questions:

1. What factors contributed to alarm fatigue in this scenario?
2. How did the telemetry optimization training improve nurse response times and patient outcomes?
3. What strategies could be implemented to sustain the improvements in telemetry management?
4. How can nurse leaders ensure ongoing adherence to telemetry discontinuation protocols?



References

Alsuyayfi, N. A., & Alanazi, A. A. (2022). Reducing non-actionable alarms in telemetry monitoring. *Journal of Clinical Nursing, 31* (5-6), 789-798. https://doi.org/10.1111/jocn.15987

Chen, Y., Wang, X., & Li, J. (2023). Impact of alarm fatigue on nurse performance in intensive care units. *Nursing in Critical Care, 28* (2), 123-130. https://doi.org/10.1111/nicc.12789

Cvach, M. (2012). Monitor alarm fatigue: An integrative review. *Biomedical Instrumentation & Technology, 46*(4), 268-277. https://doi.org/10.2345/0899-8205-46.4.268

Dee, V., Smith, R., & Jones, L. (2022). Implementing an alarm management program to reduce non-actionable alarms. *Journal of Nursing Care Quality, 37* (1), 45-52. https://doi.org/10.1097/NCQ.0000000000000598

Duffy, J. R., Culp, K., & Pugh, J. (2020). Empowering nurses through education and clinical decision-making support. *Journal of Nursing Education, 59* (11), 611-617. https://doi.org/10.3928/01484834-20201020-03

Holm, A., Larsen, T., & Petersen, C. (2024). Continuous telemetry monitoring and its impact on patient anxiety and sleep patterns. *Journal of Clinical Monitoring and Computing, 38* (1), 45-53. https://doi.org/10.1007/s10877-023-00812-4

References

Lehet, K. M., McGrath, S. P., & Bailey, J. M. (2023). Clinical decision support tools to reduce alarm fatigue. *Journal of Biomedical Informatics, 135* , 104178. https://doi.org/10.1016/j.jbi.2023.104178

Perrin, K. O., & Jones, M. (2017). Nurse -managed telemetry discontinuation protocol. *Journal of Nursing Administration, 47* (9), 425-431. https://doi.org/10.1097/NNA.0000000000000518

Rodriguez, A. M., & Thompson, J. L. (2024). Implementing telemetry discontinuation protocols to reduce hospital costs. *Journal of Healthcare Management, 69* (2), 123-132. https://doi.org/10.1097/JHM.0000000000000801

Sandau, K. E., Funk, M., Auerbach, A., Barsness, G. W., Blum, K., Cvach, M., ... & Wang, P. J. (2020). Update to practice standards for electrocardiographic monitoring in hospital settings A scientific statement from the American Heart Association. *Circulation, 141* (3), e55-e70. https://doi.org/10.1161/CIR.0000000000000754

Silverstein, M. D., & Patel, M. S. (2024). Barriers to implementing telemetry discontinuation protocols. *Journal of Hospital Medicine, 19* (1), 34-41. https://doi.org/10.12788/jhm.3792

Thank You

Appendix B: Reducing Alarm Fatigue and Enhancing Patient Safety Through Nurse Education on Telemetry Optimization Pretest

Date: _____

Purpose: This questionnaire aims to assess your current knowledge level.
Confidentiality: Your responses will be kept confidential and anonymous.
Time: The questionnaire should take approximately 10 minutes to complete.
Identification Number: Please place a six-digit identification number on this test that you will remember, as you will need to use it again following the presentation. Do not use consecutive numbers. Use a unique string of numbers (e.g., 357834). Only place the number on the test, not your name or any personal information.

Six Digit Identification Number: _____

After completing the pretest, please return it to the designated collection point.

Directions: Please select the correct response to each of the following items by circling your selection. There is only one correct response for each item.

Section 1: Multiple Choice

1. What is alarm fatigue?

A. A condition where patients become tired of hearing alarms, leading to exhaustion or annoyance.
B. A type of fatigue experienced by patients in noisy environments due to the constant barrage of alarms from medical devices.
C. A situation where alarms are too quiet to be heard, leading to missing or delayed responses.
D. A state where clinicians become desensitized to alarm sounds due to their frequency.

2. Which of the following is the primary cause of alarm fatigue?

A. Lack of patient education.
B. High frequency of alarms.
C. Poor lighting in the hospital.
D. Insufficient staffing levels.

3. How does alarm fatigue impact patient safety?

A. It improves alarm response times by making healthcare professionals more efficient in recognizing and addressing them, potentially benefiting patient safety.
B. It has no significant impact on patient safety, meaning that despite the frequent occurrence of alarms, the phenomenon does not lead to any notable changes in how healthcare providers respond to alarms.
C. It leads to slower response times or missed alarms, potentially resulting in adverse patient outcomes.
D. It reduces the number of alarms in the hospital by reducing the frequency of alarms, which could be due to changes in alarm settings or the implementation of measures to minimize unnecessary alarms, thereby potentially improving the hospital environment.

4. What is one of the environmental factors that can exacerbate alarm fatigue?

A. High patient turnover can increase stress and workload for healthcare professionals.
B. Noise pollution in healthcare settings, which leads to sensory overload.
C. Limited visiting hours, which might reduce the overall noise level in the hospital.
D. Use of electronic health records, which can streamline documentation and improve efficiency.

5. What is a significant concern about alarm fatigue in healthcare settings?

A. Desensitization and reduced responsiveness to alarms.
B. Increased patient satisfaction by reducing the frequency of alarms and creating a quieter, less stressful environment for patients.
C. Improved nurse-patient communication allows for more focused and effective interactions.
D. Enhanced use of technology, improving overall efficiency and patient care.

6. What is a significant concern about the frequency of false alarms?

A. False alarms are rare and do not affect clinicians, who can continue their duties without disruption.
B. Many false alarms can overwhelm healthcare providers, leading to alarm fatigue.
C. False alarms benefit training, allowing healthcare professionals to practice responding to alarms in a controlled environment.
D. False alarms do not affect patient safety and do not lead to significant consequences for patient care.

7. Which of the following is NOT a cause of alarm fatigue?

A. Consistent and clear alarm protocols.
B. Lack of standardization in alarm settings.
C. High frequency of alarms.
D. Environmental noise pollution.

8. What is one of the consequences of alarm fatigue mentioned in the content?

A. Delayed response to critical alarms.
B. Increased patient engagement and satisfaction.
C. Improved nurse morale and well-being.
D. Reduced need for telemetry monitoring.

9. How can alarm fatigue contribute to increased stress among healthcare providers?

A. By reducing the number of alarms they need to respond to.
B. By creating alertness and anxiety due to frequent alarms.
C. By improving their ability to manage patient care.
D. By providing more opportunities for professional development.

10. What is the primary goal of addressing alarm fatigue in healthcare settings?

A. To increase the number of alarms by flagging all potential issues.
B. To improve patient safety and clinical outcomes.
C. To reduce the use of technology in patient care.
D. To enhance patient satisfaction with hospital services.

11. According to the American Heart Association (AHA) guidelines, what are the key criteria for initiating telemetry monitoring?

A. Patient request, which implies that telemetry monitoring can be initiated based on the patient's desire to have continuous cardiac monitoring, regardless of their clinical condition or the presence of specific indications.
B. Physician preference indicates that the decision to initiate telemetry monitoring is based on the individual physician's judgment and preference rather than standardized clinical guidelines or specific patient conditions.
C. Availability of telemetry units suggests that the initiation of telemetry monitoring depends on whether telemetry units are available in the healthcare facility.
D. According to the AHA guidelines, specific cardiac conditions and clinical indications are key to initiating telemetry monitoring.

12. How does appropriate telemetry utilization impact patient safety and healthcare costs?

A. It has no impact on patient safety or costs, suggesting that the appropriate use of telemetry does not influence patient safety or the financial aspects of healthcare.
B. It increases healthcare costs, indicating that the proper utilization of telemetry leads to higher expenses for healthcare facilities.
C. It improves patient safety and reduces healthcare costs.
D. It only improves patient safety without affecting costs, implying that the correct use of telemetry enhances the safety and well-being of patients by providing continuous cardiac monitoring and timely detection of critical events.

13. What are the common causes of alarm fatigue in telemetry monitoring, and how can they be mitigated?

A. High frequency of alarms and lack of standardization; mitigated by alarm management training.
B. Low frequency of alarms and consistent protocols; mitigated by increasing alarm volume, which ensures that alarms are loud enough to be heard by healthcare providers, even in noisy environments.
C. Patient non-compliance and environmental noise; mitigated by patient education, which involves informing patients about the importance of adhering to monitoring protocols and minimizing environmental noise.
D. Limited staff training and outdated equipment; mitigated by hiring more staff, which can help distribute the workload more evenly and ensure that there are enough trained personnel to respond to alarms promptly.

14. What roles do clinical decision support tools play in optimizing telemetry utilization?

A. They replace the need for clinical judgment.
B. They guide adherence to standards.
C. They increase the number of alarms.
D. They reduce the need for telemetry monitoring.

15. How can training programs for nurses on alarm management reduce alarm fatigue and improve telemetry utilization?

A. By increasing the number of alarms.
B. By teaching nurses to ignore non-critical alarms.
C. By providing strategies to manage alarms.
D. By reducing the need for telemetry monitoring.

16. What are the benefits of adhering to AHA telemetry guidelines in the inpatient setting?

A. Increased alarm frequency and patient dissatisfaction.
B. Reduced alarm frequency and improved patient outcomes.
C. Higher healthcare costs and longer hospital stays.
D. Increased workload for nurses and more frequent alarms.

17. How does implementing a nurse-managed telemetry discontinuation protocol affect monitoring duration and alarm frequency?

A. It reduces monitoring duration and alarm frequency.
B. It does not affect monitoring duration or alarm frequency.
C. It increases monitoring duration and alarm frequency.
D. It only affects monitoring duration without changing alarm frequency.

18. What are intensive care unit nurses' training needs regarding standard cardiac monitoring practices?

A. Education on patient communication skills.
B. Education on the use of electronic health records.
C. Education on hospital administration.

D. Education on standard practices.

19. How can telemetry discontinuation education for nurse practitioners decrease hospital costs?

A. By increasing the use of continuous cardiac monitoring.
B. By reducing unnecessary monitoring.
C. By hiring more Advanced Practice Providers.
D. By extending the duration of telemetry monitoring.

20. What are the determinants of medical equipment alarm fatigue in practicing nurses, and how can these be addressed?

A. High patient turnover and limited staff; addressed by increasing staff.
B. Frequent false alarms and alarm management training.
C. Poor lighting and noise; addressed by improving hospital infrastructure.
D. Limited patient education and outdated equipment; addressed by updating equipment.

Section 2: Intentions: Please indicate your level of agreement with the following statements using the scale below:

1 = Strongly Disagree
2 = Disagree
3 = Neutral
4 = Agree
5 = Strongly Agree

Item	5 Strongly Agree	4 Agree	3 Neutral	2 Disagree	1 Strongly Disagree
1. I intend to follow the AHA guidelines for telemetry discontinuation in my daily practice.					
2. I feel confident in my ability to determine when telemetry monitoring is no longer necessary for a patient.					
3. I plan to regularly review and adhere to the telemetry discontinuation protocols provided by my institution.					

Item	5 Strongly Agree	4 Agree	3 Neutral	2 Disagree	1 Strongly Disagree
4. I believe that following the telemetry discontinuation protocol will improve patient outcomes.					
5. Based on the guidelines provided, I feel prepared to decide about discontinuing telemetry.					
6. I believe that following the telemetry discontinuation protocol will reduce the number of false alarms.					
7. I am confident I can effectively communicate the need for telemetry discontinuation to the healthcare team.					
8. I am confident in my ability to identify patients who no longer require telemetry monitoring					

Appendix C: Reducing Alarm Fatigue and Enhancing Patient Safety Through Nurse Education on Telemetry Optimization Posttest

Date: _____

Purpose: This questionnaire aims to assess your current knowledge level.

Confidentiality: Your responses will be kept confidential and anonymous.

Time: The questionnaire should take approximately 10 minutes to complete.

Identification Number: Please place the same six-digit identification number on the test you used on the pretest.

Six Digit Identification Number: _____

After completing the post-test, please return it to the designated collection point.

Directions: Please select the correct response to each of the following items by circling your selection. There is only one correct response for each item.

Section 1: Multiple Choice

1. What is alarm fatigue?

A. A condition where patients become tired of hearing alarms, leading to exhaustion or annoyance.
B. A type of fatigue experienced by patients in noisy environments due to the constant barrage of alarms from medical devices.
C. A situation where alarms are too quiet to be heard, leading to missing or delayed responses.
D. A state where clinicians become desensitized to alarm sounds due to their frequency.

2. Which of the following is the primary cause of alarm fatigue?

A. Lack of patient education.
B. High frequency of alarms.
C. Poor lighting in the hospital.
D. Insufficient staffing levels.

3. How does alarm fatigue impact patient safety?

A. It improves alarm response times by making healthcare professionals more efficient in recognizing and addressing them, potentially benefiting patient safety.
B. It has no significant impact on patient safety, meaning that despite the frequent occurrence of alarms, the phenomenon does not lead to any notable changes in how healthcare providers respond to alarms.
C. It leads to slower response times or missed alarms, potentially resulting in adverse patient outcomes.
D. It reduces the number of alarms in the hospital by reducing the frequency of alarms, which could be due to changes in alarm settings or the implementation of measures to minimize unnecessary alarms, thereby potentially improving the hospital environment.

4. What is one of the environmental factors that can exacerbate alarm fatigue?

A. High patient turnover can increase stress and workload for healthcare professionals.
B. Noise pollution in healthcare settings, which leads to sensory overload.
C. Limited visiting hours, which might reduce the overall noise level in the hospital.
D. Use of electronic health records, which can streamline documentation and improve efficiency.

5. What is a significant concern about alarm fatigue in healthcare settings?

A. Desensitization and reduced responsiveness to alarms.
B. Increased patient satisfaction by reducing the frequency of alarms and creating a quieter, less stressful environment for patients.
C. Improved nurse-patient communication allows for more focused and effective interactions.
D. Enhanced use of technology, improving overall efficiency and patient care.

6. What is a significant concern about the frequency of false alarms?

A. False alarms are rare and do not affect clinicians, who can continue their duties without disruption.
B. Many false alarms can overwhelm healthcare providers, leading to alarm fatigue.
C. False alarms benefit training, allowing healthcare professionals to practice responding to alarms in a controlled environment.
D. False alarms do not affect patient safety and do not lead to significant consequences for patient care.

7. Which of the following is NOT a cause of alarm fatigue?

A. Consistent and clear alarm protocols.
B. Lack of standardization in alarm settings.
C. High frequency of alarms.
D. Environmental noise pollution.

8. What is one of the consequences of alarm fatigue mentioned in the content?

A. Delayed response to critical alarms.
B. Increased patient engagement and satisfaction.
C. Improved nurse morale and well-being.
D. Reduced need for telemetry monitoring.

9. How can alarm fatigue contribute to increased stress among healthcare providers?

A. By reducing the number of alarms they need to respond to.
B. By creating alertness and anxiety due to frequent alarms.
C. By improving their ability to manage patient care.
D. By providing more opportunities for professional development.

10. What is the primary goal of addressing alarm fatigue in healthcare settings?

A. To increase the number of alarms by flagging all potential issues.
B. To improve patient safety and clinical outcomes.
C. To reduce the use of technology in patient care.
D. To enhance patient satisfaction with hospital services.

11. According to the American Heart Association (AHA) guidelines, what are the key criteria for initiating telemetry monitoring?

A. Patient request, which implies that telemetry monitoring can be initiated based on the patient's desire to have continuous cardiac monitoring, regardless of their clinical condition or the presence of specific indications.
B. Physician preference indicates that the decision to initiate telemetry monitoring is based on the individual physician's judgment and preference rather than standardized clinical guidelines or specific patient conditions.
C. Availability of telemetry units suggests that the initiation of telemetry monitoring depends on whether telemetry units are available in the healthcare facility.
D. According to the AHA guidelines, specific cardiac conditions and clinical indications are key to initiating telemetry monitoring.

12. How does appropriate telemetry utilization impact patient safety and healthcare costs?

A. It has no impact on patient safety or costs, suggesting that the appropriate use of telemetry does not influence patient safety or the financial aspects of healthcare.
B. It increases healthcare costs, indicating that the proper utilization of telemetry leads to higher expenses for healthcare facilities.
C. It improves patient safety and reduces healthcare costs.
D. It only improves patient safety without affecting costs, implying that the correct use of telemetry enhances the safety and well-being of patients by providing continuous cardiac monitoring and timely detection of critical events.

13. What are the common causes of alarm fatigue in telemetry monitoring, and how can they be mitigated?

A. High frequency of alarms and lack of standardization; mitigated by alarm management training.
B. Low frequency of alarms and consistent protocols; mitigated by increasing alarm volume, which ensures that alarms are loud enough to be heard by healthcare providers, even in noisy environments.
C. Patient non-compliance and environmental noise; mitigated by patient education, which involves informing patients about the importance of adhering to monitoring protocols and minimizing environmental noise.
D. Limited staff training and outdated equipment; mitigated by hiring more staff, which can help distribute the workload more evenly and ensure that there are enough trained personnel to respond to alarms promptly.

14. What roles do clinical decision support tools play in optimizing telemetry utilization?

A. They replace the need for clinical judgment.
B. They guide adherence to standards.
C. They increase the number of alarms.
D. They reduce the need for telemetry monitoring.

15. How can training programs for nurses on alarm management reduce alarm fatigue and improve telemetry utilization?

A. By increasing the number of alarms.
B. By teaching nurses to ignore non-critical alarms.
C. By providing strategies to manage alarms.
D. By reducing the need for telemetry monitoring.

16. What are the benefits of adhering to AHA telemetry guidelines in the inpatient setting?

A. Increased alarm frequency and patient dissatisfaction.
B. Reduced alarm frequency and improved patient outcomes.
C. Higher healthcare costs and longer hospital stays.
D. Increased workload for nurses and more frequent alarms.

17. How does implementing a nurse-managed telemetry discontinuation protocol affect monitoring duration and alarm frequency?

A. It reduces monitoring duration and alarm frequency.
B. It does not affect monitoring duration or alarm frequency.
C. It increases monitoring duration and alarm frequency.
D. It only affects monitoring duration without changing alarm frequency.

18. What are intensive care unit nurses' training needs regarding standard cardiac monitoring practices?

A. Education on patient communication skills.
B. Education on the use of electronic health records.
C. Education on hospital administration.
D. Education on standard practices.

19. How can telemetry discontinuation education for nurse practitioners decrease hospital costs?

A. By increasing the use of continuous cardiac monitoring.
B. By reducing unnecessary monitoring.
C. By hiring more Advanced Practice Providers.
D. By extending the duration of telemetry monitoring.

20. What are the determinants of medical equipment alarm fatigue in practicing nurses, and how can these be addressed?

A. High patient turnover and limited staff; addressed by increasing staff.
B. Frequent false alarms and alarm management training.
C. Poor lighting and noise; addressed by improving hospital infrastructure.
D. Limited patient education and outdated equipment; addressed by updating equipment.

Section 2: Intentions: Please indicate your level of agreement with the following statements using the scale below:

1 = Strongly Disagree
2 = Disagree
3 = Neutral
4 = Agree
5 = Strongly Agree

Item	5 Strongly Agree	4 Agree	3 Neutral	2 Disagree	1 Strongly Disagree
1. I intend to follow the AHA guidelines for telemetry discontinuation in my daily practice.					
2. I feel confident in my ability to determine when telemetry monitoring is no longer necessary for a patient.					
3. I plan to regularly review and adhere to the telemetry discontinuation protocols provided by my institution.					

Item	5 Strongly Agree	4 Agree	3 Neutral	2 Disagree	1 Strongly Disagree
4. I believe that following the telemetry discontinuation protocol will improve patient outcomes.					
5. Based on the guidelines provided, I feel prepared to decide about discontinuing telemetry.					
6. I believe that following the telemetry discontinuation protocol will reduce the number of false alarms.					
7. I am confident I can effectively communicate the need for telemetry discontinuation to the healthcare team.					
8. I am confident in my ability to identify patients who no longer require telemetry monitoring					

Appendix D: End-of-Session Survey

Reducing Alarm Fatigue and Enhancing Patient Safety Through Nurse Education on Telemetry Optimization

End-of-Session Survey: Educational Effectiveness

Instructions: Please take a few minutes to complete this survey. Your feedback is valuable and will help us improve future educational sessions.

1. Overall, how satisfied are you with the educational session?
   A. Very Satisfied
   B. Satisfied
   C. Neutral
   D. Dissatisfied
   E. Very Dissatisfied

2. Did the session meet your expectations? Yes No

3. Was the information presented clear and understandable? Yes No

4. Was the content relevant to your daily practice? Yes No

5. Were the objectives met? Yes No

What did you find most valuable about the session? _____

What aspects of the session could be improved? _____

Do you have any additional comments or suggestions? _____

Thank you!

Appendix E: 30-Day Post-Project Survey: Evaluating the Impact of Telemetry Optimization

Reducing Alarm Fatigue and Enhancing Patient Safety Through Nurse Education on Telemetry Optimization

30-Day Post-Project Survey: Evaluating the Impact of Telemetry Optimization

Instructions: Please take a few minutes to complete this survey. Please indicate your level of agreement with the following statements using the scale below:

Item	5 Strongly Agree	4 Agree	3 Neutral	2 Disagree	1 Strongly Disagree
9. I am comfortable applying the AHA guidelines for discontinuing telemetry in my daily practice.					
10. I feel confident in my ability to determine when telemetry monitoring is no longer necessary for a patient.					
11. To what extent do you agree that the project has successfully reduced alarm fatigue.					
12. The project positively impacted patient safety by reducing false alarms and enhancing telemetry transitions.					
13. The project has directly contributed to improving patient safety.					

1 = Strongly Disagree, 2 = Disagree, 3 = Neutral, 4 = Agree, 5 = Strongly Agree

14. Have you noticed a reduction in the number of false alarms since completing the education program? (circle one) **Yes No**

Thank You.